

MEMED BV DISTINGUISHES BETWEEN VIRAL AND BACTERIAL INFECTION IN SEPSIS PATIENTS

Einav Simon Ph.D.¹, Cesar A. Arias M.D., Ph.D.,² Richard G. Bachur M.D.³, Susanna Esposito M.D.⁴, Salim Halabi M.D.⁵, Sheldon L. Kaplan M.D.⁶, Adi Klein M.D.⁷, Sergey M. Motov M.D.⁸, Richard Rothman M.D., Ph.D.⁹, Leticia M. Ryan M.D., M.P.H.¹⁰, Shachaf Shiber M.D.¹¹, Tobias Tenenbaum M.D.¹², Alexandra Weissman M.D.¹³

¹MeMed Diagnostics, 5 Nahum Hath St. Tirat Carmel, Park High-Tech North, 39120, Israel; ²Division of Infectious Diseases, Department of Medicine, Houston Methodist Hospital, Scurlock Tower Suite 1540 6560 Fannin St, Houston, TX 77030; ³Boston Children’s Hospital, 300 Longwood Ave GL 140 Boston MA 02115, US; ⁴Pediatric Clinic, Pietro Barilla Children’s Hospital, Department of Medicine and Surgery, University of Parma, Parma, Italy; ⁵Carmel Medical Center, Mikhal St 7, Haifa, 3436212, Israel; ⁶Texas Children’s Hospital, Feigin Center, 1102 Bates Avenue, Houston, TX 77030, US; ⁷Hillel Yaffe Medical Center, Ha-Shalom St, Hadera 38100, Israel; ⁸Maimonides Medical Center, Emergency Medicine 965 48th Street, Brooklyn, NY 11219, US; ⁹Department of Emergency Medicine, 5801 Smith Avenue Suite 3220, Baltimore, MD 21209, US; ¹⁰Johns Hopkins University, 1800 Orleans Street, Sheikh Zayed Tower, Baltimore, MD 21209, US; ¹¹Rabin Medical Center, Zeev Jabotinsky St 39, Petah Tikva, 49100, Israel; ¹²Sana Klinikum Lichtenberg, Fanningerstraße 32, 10365 Berlin, Germany; ¹³University of Pittsburgh Medical Center, 3550 Terrace Street, Pittsburgh PA 15261, US

Background:

Sepsis is a life-threatening organ dysfunction syndrome caused by the body’s response to infection. Timely and appropriate sepsis management, including appropriate treatment of bacterial infection, improves outcomes. MeMed BV® (BV), a test for differentiating between bacterial and viral infection, is based on computational integration of the circulating levels of three proteins (TRAIL, IP-10, CRP). Here we evaluate its ability to differentiate bacterial from viral infection in sepsis patients.

Methods:

This was a sub-analysis of sepsis patients recruited prospectively in the Apollo study (NCT04690569). Apollo eligibility required the attending physician’s clinical suspicion of acute infection and reported fever. Sepsis was defined as two or more SIRS criteria and a suspected bacterial or viral infection classified by expert adjudication. A bacterial or viral classification required at least 2/3 experts to assign the same etiology label with confidence ≥90% or all 3 assign with confidence ≥70%. BV was measured using a platform generating a bacterial likelihood score (0-100). Based on pre-defined thresholds, scores 0-34 indicated viral (or other non-bacterial) infection, scores 35 to 65 were equivocal and 66-100 indicated bacterial infection (or co-infection). BV performance was assessed against expert panel classifications.


Conclusions:

BV accurately distinguished bacterial from viral infection in sepsis patients. This new triage tool has potential to help with timely identification of bacterial infection, enabling prompt treatment.

Results:

Seventy-nine out of 1016 eligible Apollo patients had missing heart rate and respiration rate data and a further 136 could not be classified by the experts. Out of the remaining 801 patients, 217 adults with median age of 41.8 years (IQR: 29.2-61) and 149 children with median age of 2.4 years (IQR: 1.4-5.4) had 2 or more SIRS criteria. 119 patients had at least 3 SIRS criteria and 39.6% (145/366) of the patients were hospitalized with a median duration of 4 days (IQR: 3-6 days). In the sepsis cohort, 91 patients were classified as bacterial and 275 as viral. BV yielded sensitivity and specificity of 98.8% (95%CI: 93.6-100) and 89.7% (95%CI: 85.3-93.2) and NPV of 99.6% (95%CI: 97-99.9), outperforming PCT (cut-off 0.5 ng/mL; sensitivity 52.8% (95%CI: 42-63.3); specificity 86.2% (95%CI: 81.5-90); NPV 84.6% (95%CI: 81.5-87.3)).

Patient enrollment flow



BV outperforms PCT (n = 366)

